

On the Evolution of the Temperature–Viral Mass Relation for Clusters of Galaxies

G. Mark Voit¹ & Megan Donahue²

Space Telescope Science Institute

3700 San Martin Drive

Baltimore, MD 21218

ABSTRACT

X-ray temperature measurements of clusters of galaxies are now reaching to redshifts high enough to constrain Ω_0 . A redshift-dependent relation that maps these X-ray temperatures to the virial masses of clusters is an essential ingredient when one is trying to determine cosmological parameters from cluster evolution. Most such relations assume that clusters form from top-hat perturbations that virialized just before the time we are observing them. The smaller Ω_0 is, the less accurate these relations become. Here we derive a relation between virial mass and cluster temperature that allows for the fact that clusters form gradually and cease forming when the density of the universe drops well below the critical value. We show how sensitively the expected redshift distribution of clusters depends on the mass-temperature relation used and argue that one needs to use a relation that yields no evolution in the low- Ω_0 limit.

Subject headings: cosmology: observations — galaxies: clusters: general — X-rays: galaxies

1. Introduction

Studies of temperature evolution in massive clusters of galaxies are now realizing their promise as cosmological indicators. The most massive clusters in the universe mark the

¹voit@stsci.edu

²donahue@stsci.edu

rarest, highest amplitude peaks in the initial perturbation spectrum that seeded structure formation and provide powerful leverage on parameters such as Ω_0 , the current matter density of the universe in units of the critical density ρ_{cr} . If $\Omega_0 = 1$, the development of structure should have proceeded unabated from early in time to the present day. However, the relative lack of evolution observed in the X-ray temperature function of clusters indicates that the formation of structure is stalling, and thus that $\Omega_0 \approx 0.2 - 0.5$ (Henry 1997; Bahcall et al. 1997; Carlberg et al. 1997; Donahue et al. 1998)

Using X-ray observations of clusters to constrain cosmological models requires knowledge of how the X-ray temperature of a cluster (T_X) relates to its mass. Temperatures of distant clusters are now straightforward to measure (e.g., Donahue 1997; Donahue et al. 1998; Henry 1997), but relating these temperatures to cluster masses is a subtler art. In order to define a cluster’s mass, one must decide upon an appropriate boundary separating the cluster itself from peripheral material not yet part of the cluster. Theorists tend to choose a boundary that encompasses matter that has virialized and excludes matter that is still falling into the cluster. The mass within this boundary is then M_{vir} , the virial mass of the cluster.

Defining a cluster’s mass to be M_{vir} has an established theoretical pedigree, but this definition is observationally problematic. Virial masses are a staple of the Press-Schechter approach to the problem of hierarchical structure formation, which has become enormously popular owing to its basic agreement with much more sophisticated n -body techniques (e.g., Lacey & Cole 1994). Press & Schechter (1974) presumed that the collapse of a spherical top-hat perturbation would adequately represent the formation of more complicated objects. In such a spherically symmetric collapse, the density of a newly formed object just after collapse is $18\pi^2\rho_{\text{cr}} \approx 178\rho_{\text{cr}}$ in an $\Omega_0 = 1$ universe, but this contrast factor drops to as low as $8\pi^2\rho_{\text{cr}} \approx 79\rho_{\text{cr}}$ in an open universe with $\Omega_0 \ll 1$ (e.g., Lacey & Cole 1993). Even if one could measure such density contrasts accurately, *a priori* knowledge of Ω_0 would still be needed to determine M_{vir} . Thus, the bridge between T_X and M_{vir} must be built primarily from the theoretical side.

Numerical models of cluster formation show that cluster temperatures should indeed be closely related to their masses (e.g., Evrard et al. 1996). To lowest order, cluster potentials that arise in such models are isothermal, with $T_X \propto M(r)/r$ (e.g., Cole & Lacey 1996). If the mean density within radius r is Δ times the critical density, then the mass within that radius is $M(\Delta) \propto T_X^{3/2} \rho_{\text{cr}}^{-1/2} \Delta^{-1/2}$. Proceeding in the vein of the Press-Schechter approach, one can define Δ_{vir} to be the density contrast of a spherical top-hat perturbation just after collapse and virialization, in which case $M_{\text{vir}} = M(\Delta_{\text{vir}})$ and $T_X \propto M_{\text{vir}}^{2/3} \rho_{\text{cr}}^{1/3} \Delta_{\text{vir}}^{1/3}$. The density contrast parameter Δ_{vir} depends in general on Ω_0 , the redshift of virialization, and

the value of the cosmological constant (e.g., Eke et al. 1996; Oukbir & Blanchard 1997). Adopting the recent-formation approximation, which assumes that the clusters we now observe virialized just before their light departed, then leads to a relation like the following:

$$kT_X = (1.38 \text{ keV})\beta^{-1} \left(\frac{M_{\text{vir}}}{10^{15} h^{-1} M_{\odot}} \right)^{2/3} \left[\frac{\Omega_0}{\Omega(z)} \right]^{1/3} \Delta_{\text{vir}}^{1/3} (1+z) , \quad (1)$$

where $\Omega(z)$ is the matter density of the universe in units of ρ_{cr} at redshift z , β is the usual ratio of potential depth to X-ray temperature, and h is the Hubble constant in units of $100 \text{ km s}^{-1} \text{ Mpc}^{-1}$ (e.g., Eke et al. 1996). In the limit of $\Omega(z) \approx 1$, we arrive at the scaling relation $T_X \propto M_{\text{vir}}^{2/3} (1+z)$.

The shortcomings of this approach are well known (e.g. Viana & Liddle 1996; Kitayama & Suto 1996; Eke et al. 1996). Real clusters do not form at a single moment but rather accrete their matter over a long period of time. If the accretion rate remains sufficiently high, as would be the case if $\Omega_0 = 1$, then the clusters we now see attained their observed masses relatively recently, validating the recent-formation approximation that led to equation (1). If instead $\Omega_0 \ll 1$, then the recent formation approximation grows increasingly less valid in time. Note that in the limit $\Omega_0 \rightarrow 0$, cluster evolution ceases by $z = 0$ and the $M_{\text{vir}}-T_X$ relation should no longer depend on redshift, whereas equation (1) dictates $T_X \propto M_{\text{vir}}^{2/3} (1+z)^{2/3}$. One should also keep in mind that the temperature of a real cluster depends on formation epoch only insofar as the formation epoch affects the shape of the halo density profile and furthermore that real clusters are not necessarily isothermal. Comparisons of simulated clusters with equation (1) indicate that the effective value of β declines modestly but systematically with time in an open universe with $\Omega_0 = 0.3$, slightly diminishing the expected amount of temperature evolution (Eke, Navarro, & Frenk 1998).

While the relation in equation (1) is adequate for many purposes, one would prefer to use a relation valid in the low- Ω_0 limit when constraining Ω_0 through cluster temperature evolution. One way to patch the recent-formation approximation is to define an explicit formation redshift z_f at which a cluster virializes. One can then integrate over the appropriate distribution of formation redshifts to determine the properties of observed clusters at redshift z (Kitayama & Suto 1996; Viana & Liddle 1996).

Motivated by the fact that clusters form gradually, not instantaneously, we present here an alternative approach to characterizing the redshift evolution of the $M_{\text{vir}}-T_X$ relation which has the advantage of being qualitatively correct in the low- Ω_0 limit. In § 2 we idealize clusters as forming from spherically symmetric perturbations with negative radial density gradients rather than from top-hat perturbations. This approximation alleviates the need to specify an artificial formation redshift because it explicitly accounts for how clusters grow with time. It also yields an analytical formula for the change in virial energy with

virial mass. We then use the merging-halo formalism of Lacey & Cole (1993) to derive an expression for the time-averaged mass accretion rate of a massive cluster, which allows us to determine how the $M_{\text{vir}}-T_X$ relation should change with time. Section 3 briefly illustrates some implications of this relation.

2. Mass-Temperature Relation in an Open Universe

Here we derive in three steps how the $M_{\text{vir}}-T_X$ relation should evolve. First we analyze how a radially stratified, spherically symmetric perturbation would virialize. Then we determine how the mass accretion rate of a massive cluster changes with time in a Gaussian density field with a power-law perturbation spectrum. Combining the results of these analyses, we determine how the $M_{\text{vir}}-T_X$ relation evolves with time, and we normalize this relation using equation (1). For brevity and clarity, we restrict our derivation to the analytically tractable case of an open universe without a cosmological constant.

2.1. Spherical Collapse with a Density Gradient

Let us idealize the perturbation that will form a cluster as having a spherically symmetric density profile that declines with radius r . We then can define the Lagrangian coordinate M to be the mass inside a given spherical shell centered on the origin. The evolution of each shell depends only upon the mass inside of it. In the absence of a cosmological constant, these shells will obey the familiar parametric solution $r(M) = [GM(t_M/2\pi)^2]^{1/3}(1 - \cos \theta_M)$, $t = (t_M/2\pi)(\theta_M - \sin \theta_M)$, where θ_M parametrizes the evolution of the shell containing mass M , which formally recollapses to the origin at time t_M (e.g., Peebles 1993).

Drawing an analogy between top-hat collapse and the collapse of concentric shells, we can approximate the virial mass of the forming cluster to be the M of the shell that has just recollapsed to the origin. The energy this infalling shell contributes to the total is equal to its potential energy at maximum expansion. Because the maximum radius of the shell that converges upon the origin at time t is $2[GM(t/2\pi)^2]^{1/3}$, the virial energy $-E$ of the cluster accumulates according to

$$\frac{dE}{dM_{\text{vir}}} = \frac{1}{2} \left(\frac{2\pi GM_{\text{vir}}}{t} \right)^{2/3}. \quad (2)$$

An expression for $M_{\text{vir}}(t)$ then provides the information needed to determine $E(t)$ and $T(t) \propto E(t)/M_{\text{vir}}(t)$.

2.2. Mass Accretion Rate for Massive Clusters

The merging-halo formalism of Lacey & Cole (1993) can be used to derive a mean mass accretion rate for massive clusters. This approach extends the Press-Schechter picture by considering how clusters grow via accretion of smaller virialized objects. Mass accretion becomes quasi-continuous for very massive clusters, and in this limit the Lacey & Cole formalism yields a mass accretion rate that enables us to integrate equation 2.

In order to analyze halo merger rates, one defines $\delta(\mathbf{x}, t; M)$ to be the local fractional overdensity of the universe, smoothed on mass scale M and centered on comoving point $\mathbf{x}$ at time t . Early in time the perturbations described by δ are assumed to be Gaussian, and in the linear regime they grow in proportion to the function $D(t)$, which depends on Ω_0 . While these perturbations remain linear, their rms amplitude on scale M can be expressed as $\sigma(M)D(t)/D(t_0)$. Ultimately some of them grow non-linear, and they are assumed to virialize when their amplitudes, extrapolated from the linear regime according to $D(t)$, exceed some critical threshold $\delta_c(t)$. One can then trace the merger history of a mass parcel beginning at $\mathbf{x}$ from time t_1 to the present by keeping track of the largest M for which $\delta(\mathbf{x}, t_1; M)D(t)/D(t_1) > \delta_c(t)$. This largest mass is the mass of the virialized halo containing the mass parcel at time t , and it jumps by an amount ΔM each time the halo merges with another of mass ΔM .

Equivalently, one can analyze the growth of structure by keeping the amplitudes of the perturbations fixed, with rms amplitude $\sigma(M)$, and tracking when they exceed the steadily falling threshold $\omega(t) \equiv \delta_c(t)D(t_0)/D(t)$. In an open universe with no cosmological constant, Lacey & Cole (1993) show that

$$\omega(t) = \frac{3}{2}D(t_0)[1 + (t_\Omega/t)^{2/3}] \quad (3)$$

where $t_\Omega = (\pi\Omega_0/H_0)(1 - \Omega_0)^{-3/2}$. The parameter ω thus serves as a useful surrogate for the time coordinate in an open universe because it asymptotically approaches a fixed value as growth of structure slows to a halt.

Now we are ready to determine an expression for the halo accretion rate. Lacey & Cole (1993) show that the probability that the mass of a virialized halo will jump from M_1 into an interval dM_2 at M_2 within a time interval corresponding to $d\omega$ is

$$\frac{d^2p}{dS_2d\omega}(S_1 \rightarrow S_2|\omega)dS_2d\omega = \frac{1}{(2\pi)^{1/2}} \left[\frac{S_1}{S_2(S_1 - S_2)} \right]^{3/2} \exp \left[-\frac{\omega^2(S_1 - S_2)}{2S_1S_2} \right] dS_2d\omega \quad , \quad (4)$$

where $S_i \equiv \sigma^2(M_i)$. To find the mean growth rate of a halo of mass M , we can integrate

over this probability distribution as follows:

$$\left\langle \frac{dS}{d\omega} \right\rangle = \int_{S_2}^{\infty} (S_1 - S_2) \frac{d^2 p}{dS_2 d\omega} dS_1 . \quad (5)$$

The variance $S(M)$ generally declines with M , so for a rare high-mass object, $S_2(M_2) \ll \omega^2$. In this limit, the integrand's exponential behavior restricts its main contribution to a narrow range where $S_1 \approx S_2$, reflecting the fact that most of the accreted mass comes from objects much smaller than the cluster. Thus,

$$\left\langle \frac{dS}{d\omega} \right\rangle \approx \frac{S}{\omega} . \quad (6)$$

If the fluctuation amplitudes obey $\sigma(M) \propto M^{-(n+3)/6}$, where n is the usual power-law perturbation index in wavenumber space, we finally arrive at the mass accretion law $M_{\text{vir}} \propto \omega^{-3/(n+3)}$. This result is equivalent to Lacey & Cole's (1993) finding that $M_{\text{vir}} \propto t^{2/(n+3)}$ for $\Omega_0 = 1$.

2.3. Evolution and Normalization of $M_{\text{vir}}-T_X$

The typical temperature of a cluster should be proportional to the virial energy divided by M_{vir} . If we define $x \equiv [1 + (t_{\Omega}/t)^{2/3}]$ and $m \equiv 5/(n+3)$, we can write $M_{\text{vir}} \propto x^{-3m/5}$ and $dE \propto (x-1)dx^{-m}$. Integrating over x then gives $E \propto M_{\text{vir}}^{5/3}(x-1+1/m)$. Thus, we obtain

$$T_X \propto \frac{E}{M_{\text{vir}}} \propto M_{\text{vir}}^{2/3} \left[\left(\frac{t_{\Omega}}{t} \right)^{2/3} + \left(\frac{n+3}{5} \right) \right] . \quad (7)$$

This relation behaves properly in all the appropriate limits. The time-dependent factor varies like $(1+z)$ when $\Omega(z) \approx 1$, like $\rho_{\text{cr}}^{1/3} \propto t^{-2/3}$ for intermediate $\Omega(z)$, and goes to a constant as $\Omega(z) \rightarrow 0$. To normalize this expression, we can simply match it to equation (1) at early times:

$$kT_X = (2.76 \text{ keV})\beta^{-1} \left(\frac{M_{\text{vir}}}{10^{15} h^{-1} M_{\odot}} \right)^{2/3} \frac{1 - \Omega_0}{\Omega_0^{2/3}} \left[\left(\frac{t_{\Omega}}{t} \right)^{2/3} + \left(\frac{n+3}{5} \right) \right] . \quad (8)$$

Reducing Ω_0 at fixed M_{vir} and t pushes T_X upward because the effective formation epoch moves to earlier times, when the universe was denser. Note also that the implied asymptotic behavior of temperature evolution depends on the power-law index n . Larger amounts of power on small scales (larger n) lead to a less sensitive dependence of M_{vir} on ω and an earlier cutoff for the accumulation of mass.

The true test of a relation between M_{vir} and T_X is how accurately it reproduces the evolution of the cluster temperature function in a large numerical simulation. Showing that $M(\Delta)$ correlates well with T_X for some chosen Δ is not sufficient because such tests do not probe how Δ_{vir} changes with time. As long as clusters are nearly isothermal above the density contrast Δ , one will find $T_X \propto [M(\Delta)]^{2/3} \rho_{\text{cr}}^{1/3} \Delta^{1/3}$. If the goal is to constrain cosmological parameters by applying Press-Schechter analyses to observed clusters, then the $M_{\text{vir}}-T_X$ relation one uses should yield cluster temperature functions that approximate the results of numerical simulations as closely as possible. To accurately reproduce behavior at low Ω , one needs a relation like equation (7) that asymptotically yields no evolution. In the most general cases where a single value of n does not adequately represent the perturbation spectrum, equation (7) should still be useful as a fitting formula in which the asymptotic constant is a free parameter.

3. Press-Schechter Predictions and $M_{\text{vir}}-T_X$ Evolution

The generally good agreement between analytical Press-Schechter models of cluster evolution and numerical simulations has inspired numerous applications of the Press-Schechter approach to X-ray cluster surveys (e.g., Oukbir & Blanchard 1997; Henry 1997; Carlberg et al. 1997; Donahue et al. 1998). An accurate $M_{\text{vir}}-T_X$ relation is a crucial element in such studies because it maps the virial masses used in the Press-Schechter formalism to either observed X-ray temperatures or observed X-ray luminosities (L_X) through an L_X-T_X relation. Here we will show that the consequences of using an inaccurate $M_{\text{vir}}-T_X$ relation can be quite significant.

For the purposes of studying cluster evolution, the change in $M_{\text{vir}}-T_X$ with z is more important than its normalization. Figure 1 shows how the quantity $T_X(M_{\text{vir}}, z)/T_X(M_{\text{vir}}, 0)$ changes for four different relations: the one derived here for $n = -1$ and -2 , the relation from equation (1), and the relation $T_X \propto M_{\text{vir}}^{2/3}(1+z)$ that one obtains by assuming that the mean density within the virial radius is some constant multiple of the background density. We have also computed the mean temperatures for clusters of a fixed mass at a variety of redshifts by integrating over formation redshifts following Viana & Liddle’s (1996) method and with Lacey & Cole’s (1993) definition of formation epoch. The points on Figure 1 show how the mean temperatures of these clusters that are originally 3σ perturbations evolve for $n = -1$. Note the close agreement between these points and the $n = -1$ $M_{\text{vir}}-T_X$ relation from the present paper.

The differences between the $M_{\text{vir}}-T_X$ relations in Figure 1 are relatively modest in percentage terms, but they become greatly amplified when filtered through the exponential

distribution of cluster masses that emerges from Press-Schechter calculations. According to such calculations, the differential comoving number density of clusters dn in temperature interval dT_X is given by

$$\frac{dn}{dT_X} = \left(\frac{2}{\pi}\right)^{1/2} \frac{3}{2} \frac{\Omega_0 \rho_{\text{cr}}(0)}{T_X M_{\text{vir}}(T_X, z)} \frac{d \ln \sigma}{d \ln M_{\text{vir}}} \nu_c(T_X, z) \exp[-\nu_c^2(T_X, z)/2] . \quad (9)$$

For a power-law perturbation index n and a fiducial temperature T_0 , one can write $\nu_c(T_X, z) = \nu_{c0}[\omega(z)/\omega(0)][M_{\text{vir}}(T_X, z)/M_{\text{vir}}(T_0)]^{(n+3)/6}$, where ν_{c0} or some analogous parameter is adjusted to fit low-redshift cluster surveys: fixing ν_{c0} is equivalent to fixing the amplitude of the power spectrum for a given n . The derived number density of clusters thus depends exponentially on $M_{\text{vir}}(T_X, z)$.

An $M_{\text{vir}}-T_X$ relation that overestimates temperature evolution will underestimate the virial mass corresponding to a high- z cluster of a given temperature or luminosity. Because the number density of clusters above a given mass decreases with virial mass, an underestimate of virial mass leads one to *overestimate* the numbers of high- z clusters that ought to be in a particular temperature-limited or flux-limited sample. This effect is more severe in a flux-limited sample because the expected luminosity is much more sensitive to changes in the virial mass.

To illustrate these effects, we have computed the surface density of clusters on the sky given by equation (9) for $\Omega_0 = 0.2$, $n = -1$, various $M_{\text{vir}}-T_X$ relations, the L_X-T_X relation from Edge & Stewart (1991), which we assume remains constant with redshift (e.g., Mushotzky & Scharf 1997), and a particular set of selection criteria. We restrict the clusters to have $T_X > 5 \text{ keV}$ and $F_X > 10^{13} \text{ erg cm}^{-2} \text{ s}^{-1}$ within a $2.4' \times 2.4'$ detection cell like that of the Einstein Extended Medium Sensitivity Survey (EMSS; Henry et al. 1992). We also include the effects of Galactic soft X-ray absorption, assuming a mean hydrogen column density $3 \times 10^{20} \text{ cm}^{-2}$. Figure 2 shows the resulting surface densities, binned in redshift intervals of 0.1. Note that at $z \sim 0.5$ the expected cluster counts differ by a factor ~ 3 , with more modest temperature evolution yielding fewer expected clusters. This factor is similar to the discrepancy Oukbir & Blanchard (1997) find between the actual redshift distribution of clusters in the EMSS and their expectations, suggesting that the entire discrepancy might stem from the $M_{\text{vir}}-T_X$ relation they use. Above $z \sim 0.5$, the flux limit becomes more restrictive than the temperature limit, and the discrepancies between the expected cluster counts grow more pronounced.

Clearly, Press-Schechter approaches to modelling X-ray surveys require an accurate $M_{\text{vir}}-T_X$ relation. The relation we give here is likely to describe temperature evolution more accurately at late times than those based on the recent-formation approximation because it correctly yields no evolution as $\Omega(z) \rightarrow 0$. We are currently using this $M_{\text{vir}}-T_X$

relation to study cluster evolution in the EMSS and will report our results in a future paper.

We would like to thank Caleb Scharf for his helpful suggestions while we were developing these ideas. We acknowledge NASA grants grant NAG5-3257 and NAG5-3208 for partial support.

REFERENCES

- Bahcall, N. A., Fan, X., & Cen, R. 1997, *ApJ*, 485, L53.
- Carlberg, R. G., Morris, S. L., Yee, H. K. C., & Ellingson, E. 1997, *ApJ*, 479, L19.
- Cole, S., & Lacey, C. 1996, *MNRAS*, 281, 716
- Donahue, M. 1996, *ApJ*, 468, 79.
- Donahue, M., Voit, G. M., Gioia, I., Luppino, G., Hughes, J. P., & Stocke, J. T. 1998, *ApJ*, in press
- Edge, A., & Stewart, G. 1991, *MNRAS*, 252, 428
- Eke, V. R., Cole, S., & Frenk, C. S. 1996, *MNRAS*, 282, 263.
- Eke, V. R., Navarro, J. F., & Frenk, C. S. 1998, *ApJ*, in press
- Evrard, A. E., Metzler, C. A. & Navarro, J. F. 1996, *ApJ*, 469, 494.
- Henry, J. P., Gioia, I. M., Maccacaro, T., Morris, S. L., Stocke, J. T. 1992, *ApJ*, 386, 408.
- Henry, J. P. 1997, *ApJ*, 489, L1
- Kitayama, T., & Suto, Y. 1996, *ApJ*, 469, 480
- Lacey, C., & Cole, S. 1994, *MNRAS*, 271, 676
- Lacey, C., & Cole, S. 1993, *MNRAS*, 262, 627
- Mushotzky, R., & Scharf, C. A. 1997, *ApJ*, 482, L13.
- Oukbir, J., & Blanchard, A. 1997, *A&A*, 317, 1.
- Peebles, P. J. E. 1993, *Principles of Physical Cosmology*, Princeton: Princeton University Press.
- Press, W., & Schechter, P. 1974, *ApJ*, 187, 425.
- Viana, P. T. P., & Liddle, A. R. 1996, *MNRAS*, 281, 323.




Fig. 1.— Temperature evolution for various $M_{\text{vir}}-T_X$ relations. The naive relation in which $T_X \propto M_{\text{vir}}^{2/3}(1+z)$ (long-dashed line) predicts the most evolution in T_X for a given M_{vir} . Temperature evolution is more modest under the late-formation approximation in an open universe (dotted line) and is milder still for the relation derived in equation (7), given $n = -1$ (solid line) and $n = -2$ (short-dashed line). The points indicate how mean cluster temperatures evolve in models that integrate over a range of formation redshifts z_f to find the temperatures of clusters at a given redshift z , given an $n = -1$ perturbation spectrum.




Fig. 2.— Redshift distribution of clusters as predicted by different $M_{\text{vir}}-T_X$ relations. Here we show expectations for clusters hotter than 5 keV with fluxes exceeding $10^{13} \text{ erg cm}^{-2} \text{ s}^{-1}$ in the EMSS detect cell. The naive relation $T_X \propto M_{\text{vir}}^{2/3} (1+z)$ (dashed line) overpredicts the high- z clusters because it overestimates temperature evolution and maps the flux and temperature limits to lower values of M_{vir} than it should.